

THESES PUBLICATIONS

Adaes S, Ferreira-Gomes J, Mendonca M, Almeida L, Castro-Lopes JM, Neto FL. Injury of primary afferent neurons may contribute to osteoarthritis induced pain: an experimental study using the collagenase model in rats. *Osteoarthritis and Cartilage*. 2015;23(6):914-24.

Adaes S, Mendonca M, Santos TN, Castro-Lopes JM, Ferreira-Gomes J, Neto FL. Intra-articular injection of collagenase in the knee of rats as an alternative model to study nociception associated with osteoarthritis. *Arthritis Research & Therapy*. 2014;16(1).

Adães S., Almeida L., Potes C.S., Ferreira A.R., Castro-Lopes JM., Ferreira-Gomes J., Neto F.L. (2017) Glial activation in the collagenase model of nociception associated with osteoarthritis. *Molecular Pain* (in press)

Aguiar P, Sousa M, Szucs P. Versatile morphometric analysis and visualization of the three-dimensional structure of neurons. *Neuroinformatics* 11: 393-403 (2013)

Alemi M, Cristiana Gaiteiro, Carlos Alexandre Ribeiro, Luís Miguel Santos, João Rodrigues Gomes, Sandra Marisa Oliveira, Pierre-Olivier Couraud, Babette Weksler, Ignacio Romero, Maria João Saraiva & Isabel Cardoso. Transthyretin participates in beta amyloid transport from the brain to the liver- involvement of the low density lipoprotein receptor-related protein 1? *Scientific Reports* | 6:20164 | DOI: 10.1038/srep20164

Alemi M, Sara Costa Silva, Isabel Santana, Isabel Cardoso. Transthyretin stability is critical in assisting beta amyloid clearance - Relevance of Transthyretin stabilization in Alzheimer's Disease. *CNS Neuroscience & Therapeutics* (in press)

Angeliki M, S. Pavlou, Pinho R, Paiva I, and Fleming Outeiro T. The yin and yang of a-synuclein-associated epigenetics in Parkinson's disease, *Brain*, 2017

Baptista MV, Ferreira S, Pinho-e-Melo T, Carvalho M, Cruz VT, Carmona C, et al. Mutations of the GLA Gene in Young patients with Stroke The PORTYSTROKE Study- Screening Genetic Conditions in PORTuguese Young STROKE Patients. *Stroke*. 2010 Mar;41(3):431-6.

Borges GS, Berrocoso E, Ortega-Alvaro A, Mico JÁ, Neto FL. Extracellular signal-regulated kinase activation in the chronic constriction injury model of neuropathic pain in anaesthetized rats. *Europ. J. Pain* 17: 35-45, 2013

Borges GS, Mico JA, Neto FL, Berrocoso E. Corticotrophin-releasing factor mediates pain-induced anxiety through the ERK1/2 signalling cascade in Locus Coeruleus neurons. *Int J Neuropsychopharmacol*. 2015 Feb 25;18(8). pii: pyv019. doi: 10.1093/ijnp/pyv019.

Borges GS, Neto FL, Mico JÁ, Berrocoso E. Reversal of monoarthritis-induced affective disorders by diclofenac in rats. *Anesthesiology* 120:1476-1490 (2014).

Borges GS, Miguelez C, Neto FL, Mico JÁ, Ugedo L, Berrocoso E. Activation of extracellular signal-regulated kinases (ERK1/2) in Locus Coeruleus contributes to pain-related anxiety in monoarthritic rats. Jun 2017, 20(6):463. doi: 10.1093/ijnp/pyx005.

Branco P, DanielaSeixas, SabineDeprez, SilviaKovacs, RonaldPeeter, SãoL.Castro and StefanSunaert. Resting State Functional Magnetic Resonance Imaging for Language Preoperative Planning. *Frontiers in Human Neuroscience*, published: 01 February 2016 doi: 10.3389/fnhum.2016.00011

Branco P, Daniela Seixas, São Luís Castro. Temporal reliability of ultra-high field resting-state MRI for single-subject sensorimotor and language mapping. *NeuroImage* (e-pub ahead of print)

Cardoso A, Assunção M, Andrade JP, Pereira PA, Madeira MD, Paula-Barbosa MM, Lukoyanov NV. Loss of synapses in the entorhinal-dentate gyrus pathway following repeated induction of electroshock seizures in the rat. *Journal of Neuroscience Research* 86:71-83, 2008.

Cardoso A, Carvalho LS, Lukoyanova EA, Lukoyanov NV. Effects of repeated electroconvulsive shock seizures and pilocarpine-induced status epilepticus on emotional behavior in the rat. *Epilepsy & Behavior* 14:293-299, 2009.

Cardoso A, Lukoyanova EA, Madeira MD, Lukoyanov NV. Seizure-induced structural and functional changes in the rat hippocampal formation: Comparison between brief seizures and status epilepticus. *BEHAVIOURAL BRAIN RESEARCH* 225: 538-546 (2011)

Cardoso A, Lukoyanova EA, Madeira MD, Lukoyanov NV. Seizure-induced structural and functional changes in the rat hippocampal formation: Comparison between brief seizures and status epilepticus. *BEHAVIOURAL BRAIN RESEARCH* 225: 538-546 (2011)

Cardoso A, Madeira MD, Paula-Barbosa MM, Lukoyanov NV. Retrosplenial granular b cortex in normal and epileptic rats: A stereological study *Brain Research* 1218:206-214, 2008.

Cardoso A, Paula-Barbosa MM, Lukoyanov NV. Reduced density of neuropeptide Y neurons in the somatosensory cortex of old male and female rats: Relation to cholinergic depletion and recovery after nerve growth factor treatment. *Neuroscience* 137:937-948, 2006.

Cardoso-Cruz H*, Dourado M*, Monteiro C, Matos M, Galhardo V (2014) Activation of dopaminergic D2/D3 receptors modulates dorsoventral connectivity in the hippocampus and reverses pain-induced impairment of working memory. *The Journal of Neuroscience*, 34:5861-5873. (DOI: 10.1523/JNEUROSCI.0021-14.2014)

Cardoso-Cruz, H, Lima, D, Galhardo, V Instability of spatial encoding by CA1 hippocampal place cells after peripheral nerve injury (2011) *EUROPEAN JOURNAL OF NEUROSCIENCE* Volume: 33 Pages: 2255-2264

Cardoso-Cruz, H, Sousa, Vieira, JB, Lima, D, Galhardo, V Prefrontal cortex and mediodorsal thalamus reduced connectivity is associated with spatial working memory impairment in rats with inflammatory pain (2013) *PAIN* Volume: 154 Pages: 2397-2406

Cardoso-Cruz, H; Sameshima, Koichi; Lima, Deolinda; Galhardo, Vasco. Dynamics of Circadian Thalamocortical Flow of Information during a Peripheral Neuropathic Pain Condition. (2011) *Frontiers in integrative neuroscience* Volume:5 Pages:43

Cardoso-Cruz, H; Lima, D; Galhardo, V. Impaired Spatial Memory Performance in a Rat

Model of Neuropathic Pain Is Associated with Reduced Hippocampus–Prefrontal Cortex Connectivity (2013), The Journal of Neuroscience, 33(6):2465-80.

Chen, ZF, Rebelo, S, White, F, Malmberg, AB, Baba, H, Lima, D, Woolf, C, Basbaum, AI, Anderson, DJ The paired homeodomain protein DRG11 is required for the projection of cutaneous sensory afferent fibers to the dorsal spinal cord (2001) NEURON Volume: 31 Pages: 59-73

Coelho A, Cruz F, Cruz CD, Avelino A (2012) Effect of Onabotulinumtoxin/A on intramural parasympathetic ganglia: an experimental study on guinea pig bladder. J Urol 187(3): 1121-6

Coelho A, Cruz F, Cruz CD, Avelino A (2012) Spread of Onabotulinumtoxin/A after bladder injection. Experimental study using the distribution of cleaved SNAP-25 as the marker of the toxin action. Eur Urol 61(6):1178-1184

Coelho A, Dinis P, Pinto R, Gorgal T, Silva C, Silva A, Silva J, Cruz CD, Cruz F, Avelino A (2010) Distribution of the high-affinity binding sites and intracellular target of Botulinum Toxin type A in the Human Bladder. Eur Urol 57:884-890

Coelho A, Oliveira R, Rossetto O, Cruz CD, Cruz F, Avelino A. (2014) Intrathecal administration of botulinum toxin type A improves urinary bladder function and reduces pain in rats with cystitis. Eur J Pain. 2014 Apr 23. doi: 10.1002/ejp.513. [Epub ahead of print]

Dias C, Gaio AR, Monteiro E, Barbosa S, Cerejo A, Donnelly J, et al. Kidney-Brain Link in Traumatic Brain Injury Patients? A preliminary report. Neurocritical care. 2015;22(2):192-201.

Dias C, Maia I, Cerejo A, Varsos G, Smielewski P, Paiva JA, et al. Pressures, Flow, and Brain Oxygenation During Plateau Waves of Intracranial Pressure. Neurocritical Care. 2014;21(1):124-32.

Dias C, Rocha J, Pereira E, Cerejo A. Traumatic Brain Injury in Portugal: Trends in Hospital Admissions from 2000 to 2010. Acta Medica Portuguesa. 2014;27(3):349-56.

Dias C, Silva MJ, Pereira E, Monteiro E, Maia I, Barbosa S, et al. Optimal Cerebral Perfusion Pressure Management at Bedside: A Single-Center Pilot Study. Neurocrit Care. 2015.

Dias C, Silva MJ, Pereira E, Silva S, Cerejo A, Smielewski P, et al. Post-Traumatic Multimodal Brain Monitoring: Response to Hypertonic Saline. Journal of Neurotrauma. 2014;31(22):1872-80.

Dias C, Multimodal brain monitoring in neurocritical care practice (2014), The international Journal of Neurosciences and mental health, 1(Suppl. 1):S08.

Dourado M, Cardoso-Cruz H, Monteiro C, Galhardo V. Effect of motor impairment on analgesic efficacy of dopamine D2/3receptors in a rat model of neuropathy. Journal of Experimental Neuroscience 2016, 10:51-57

Fernandes S, Salta S, Bravo J, Silva AP, Summavielle T. Acetyl-L-Carnitine Prevents Methamphetamine-Induced Structural Damage on Endothelial Cells via ILK-Related MMP-9 Activity. Molecular neurobiology. 2014.

Fernandes S, Salta S, Summavielle T. Methamphetamine promotes alpha-tubulin deacetylation in endothelial cells: The protective role of acetyl-L-carnitine. *Toxicology Letters*. 2015;234(2):131-8.

Ferreira DG, Temido-Ferreira M, Vicente Miranda H, Batalha VL, Coelho JE, Szegö EM, Marques-Morgado I, Vaz SH, Rhee JS, Schmitz M, Zerr I, Lopes LV, and Outeiro TF (2017) α -Synuclein interacts with PrPC to induce cognitive impairment through mGluR5 and NMDAR2B. *Nature Neuroscience* (in press)

Ferreira DG, Batalha VL, Vicente Miranda H, Coelho JE, Gomes R, Gonçalves FQ, Real JI, Rino J, Albino-Teixeira A, Cunha RA, Outeiro TF, and Lopes LV (2017) Adenosine A2A Receptors Modulate α -Synuclein Aggregation and Toxicity. *Cereb Cortex* 27(1):718-730. doi: 10.1093/cercor/bhv268

Ferreira S, Auray-Blais C, Boutin M; Lavoie P, Nunes JP, Martins E, Garman S, Oliveira JP. Variations in the GLA gene correlate with globotriaosylceramide and globotriaosylsphingosine analog levels in urine and plasma. *Clin Chim Acta*. 2015 Jul 20; 447:96-104. doi: 10.1016/j.cca.2015.06.003. Epub 2015 Jun 9.

Ferreira S, Reguenga C, Oliveira JP. The modulatory effects of the polymorphisms in GLA 5'-untranslated region upon gene expression are cell-type specific. *JIMD Rep*. 2015; 23:27-34. doi: 10.1007/8904_2015_424. Epub 2015 Mar 13.

Ferreira S, Ortiz A; Germain DP, Viana-Baptista M, Caldeira Gomes A, Camprecios M, Fenollar-Cortés M, Gallegos-Villalobos Á, Garcia D, García-Robles JA Egido J, Gutiérrez-Rivas E, Herrero JÁ, Mas S, Oancea R, Péres P, Salazar-Martín LM, Solera-Garcia J, Alves H, Garman SC, Oliveira JP. The alpha-galactosidase A p. Arg118Cys variant does not cause a Fabry disease phenotype: Data from individual patients and family studies. *Mol Genet Metab*. 2015 Feb; 114(2):248-258. doi: 10.1016/j.ymgme.2014.11.004. Epub 2014 Nov 9.

Ferreira S, Viana-Baptista M, Rodrigues D, Oliveira JP. Individuals with normal GLA sequence may present abnormally spliced alpha-galactosidase mRNA transcripts. *Gene Cell and Tissue*. 2016; 3 (1):e34386

Ferreira-Gomes J, Adães S, Castro-Lopes JM. (2008) Assessment of movement-evoked pain in osteoarthritis by the Knee-bend and CatWalk tests: a clinically relevant study. *Journal of Pain* 9: 945-954

Ferreira-Gomes J, Adães S, Mendonça M, Castro-Lopes JM. (2012) Analgesic effects of lidocaine, morphine and diclofenac on movement-induced nociception, as assessed by the Knee-Bend and CatWalk tests in a rat model of osteoarthritis. *Pharmacology Biochemistry and Behavior* 101(4):617-24: a clinically relevant study. *Journal of Pain* 9: 945-954

Ferreira-Gomes J, Adães S, Sarkander J, Castro-Lopes JM. (2010) Phenotypic alterations of neurons that innervate osteoarthritis joints in rats. *Arthritis&Rheumatism* 62(12): 3677-85.

Ferreira-Gomes J, Adães S, Sousa RM, Mendonça M, Castro-Lopes JM. (2012) Dose-dependent expression of neuronal injury markers during experimental osteoarthritis induced by monoiodoacetate in the rat. *Molecular Pain* 8:50.

Frias B, Allen S, Dawbarn D, Charrua A, Cruz F, Cruz CD (2013) Brain-derived neurotrophic factor, acting at the spinal cord level, participates in bladder hype

ractivity and referred pain during chronic bladder inflammation. *Neuroscience*. 234:88-102.

Frias B, Charrua A, Avelino A, Michel MC, Cruz F, Cruz CD (2012) Transient receptor potential vanilloid 1 mediates nerve growth factor-induced bladder hyperactivity and noxious input. *BJU Int*. 110(8 Pt B): E422-8.

Frias B, Santos J, Morgado M, Sousa M, Allen S, Cruz F, Cruz CD (2013). The role of Brain Derived Neurotrophic Factor (BDNF) in the development of neurogenic detrusor overactivity (NDO). *J. Neurosci*.35: 2146-2160 (2015)

Frias, B, Lopes, T, Pinto, R, Cruz, F, Cruz, CD Neurotrophins in the Lower Urinary Tract: Becoming of Age (2011) *Current Neuropharmacology* Volume: 9 Pages: 553-558

Goadsby PJ, Holland PR, **Martins-Oliveira M**, Hoffmann J, Schankin C, Akerman S. (2017) Pathophysiology of migraine: A disorder of sensory processing. *Physiol. Rev*. 97:553-622

Gomes CP, **Lopes CDF**, Moreno PMD, Varella-Moreira A, Alonso MJ, Pêgo AP. Translating chitosan to clinical delivery of nucleic acid-based drugs. *MRS Bulletin* 2014; 39:60-70

Guimaraes, J ; Vieira-Coelho, A ; Rosas, MJ ; Moura, E ; Vaz, R; Garrett, C Acute Renal Failure in Patients with Bilateral Deep Brain Stimulation. *Mov Disord* 2010 Oct; 25(14):2462-2464.

Guimaraes, J ; Vieira-Coelho, A ; Rosas, MJ ; Moura, E ; Vaz, R; Garrett, C. Follow-up of renal function in Parkinson patients with bilateral deep brain stimulation. *Parkinsonism Relat Disord* 2013 Sep; 19(9):836-837

Guimaraes, J., Moura, E., Vieira-Coelho, M. A., & Garrett, C. (2012). Weight variation before and after surgery in Parkinson's disease: a noradrenergic modulation? [Review]. *Mov Disord*, 27(9), 1078-1082. doi: 10.1002/mds.25063

Guimaraes, J; Moura E; Elisabete Silva; Paulo Aguiar; Carolina Garrett; Maria Augusta Vieira-Coelho. Locus coeruleus is involved in weight loss in a rat model of Parkinson's disease: an effect reversed by deep brain stimulation. *BRAIN STIMULATION* 2013 Jun; 6: 845-855 DOI: 10.1016/j.brs.2013.06.002

Guimaraes, J; Matos, E; Rosas, MJ; Vieira-Coelho, A; Borges, N; Correia, F; Vaz, R; Garrett, C. Modulation of nutritional state in Parkinsonian patients with bilateral subthalamic nucleus stimulation. *J Neurol* 2009 Dec; 256(12):2072-2078.

Guimaraes, J; Moura, E; Vieira-Coelho, MA; Garrett, C. Weight variation before and after surgery in Parkinson's disease: a noradrenergic modulation? *Mov Disord* 2012 Aug; 27(9):1078-1082.

Guimaraes, J; Vieira-Coelho, MA; Moura, E; Afonso, J; Rosas, MJ; Vaz, R; Garrett, C. Urinary profile of catecholamines and metabolites in Parkinson patients with deep brain stimulation. *Eur J Neurol* 2013 May; 2: 353-356, DOI: 10.1111/ene.12161

Leite-Almeida H, **Melo A**, Pego JM, Bernardo S, Milhazes N, Borges F, et al. Variable delay-to-signal: a fast paradigm for assessment of aspects of impulsivity in rats. *Frontiers in Behavioral Neuroscience*. 2013;7.

Liu Y, Abdel Samad O, Zhang L, Duan B, Tong Q, **Lopes C**, Ji RR, Lowell BB, Ma Q. (2010) VGLUT2-dependent glutamate release from nociceptors is required to sense pain and suppress itch. *Neuron* 2010 Nov 4;68(3):543-56.

Lopes C, Liu Z, Xu Y, Ma Q (2012) Tlx3 and Runx1 act in combination to coordinate the development of a cohort of nociceptors, thermoceptors, and pruriceptors. *J Neurosci*. 2012 Jul 11;32(28):9706-15.

Lopes CDF, Gomez-Lazaro M, Pêgo AP. Seeing is believing but quantifying is deciding. *Nanomedicine* 2015, 10: 2307-2310

Lopes CDF, Gomes CP, Neto E, Sampaio P, Aguiar P, Pêgo AP. Microfluidic-based platform to mimic the in vivo peripheral administration of neurotropic nanoparticles. *Nanomedicine (Lond)*. (2016)11(24):3205-3221

Lopes CDF, Gonçalves N, Gomes CP, Saraiva MJ, Pêgo AP "BDNF gene delivery mediated by neuron-targeted nanoparticles is neuroprotective in peripheral nerve injury" *Biomaterials* (2017) 0.1016/j.biomaterials.2016.12.025, 1 (2017) 83-96

Lopes CDF, Oliveira H, Estevão I, Pires LR, Pêgo AP. In vivo targeted gene delivery to peripheral neurons mediated by neurotropic poly(ethylene-imine)-based nanoparticles. *Int J Nanomed* 2016, 11: 2675-2683

Luz LL, Szucs P, Pinho R & Safronov BV. Monosynaptic excitatory inputs to spinal lamina I anterolateral-tract-projecting neurons from neighbouring lamina I neurons. *J Physiol* 588, 4489-4505, 2010

Maia GH, Quesado JL, Soares JI, doCarmo JM, Andrade PA, Andrade JP, Lukoyanov NV. Loss of hippocampal neurons after kainate treatment correlates with behavioral deficits. *Plos ONE* 2014, 9:e84722

Maia GH, Soares JI, Andrade PA, Leite JF, Andrade JP, Lukoyanov NV. Altered taste preference and loss of limbic-projecting serotonergic neurons in the dorsal raphe nucleus of chronically epileptic rats. *Behav Brain Res* 297: 28-38 (2016)

Maia GH, Brazete CS, Soares JI, Luz LL, Lukoyanov NV. Serotonin depletion increases seizure susceptibility, facilitates epileptogenesis and worsens neurological outcome in kainate model of epilepsy. *Brain Research Bulletin*, 2016 Jan 15;297:28-36.

Marques-Lopes, J, Martins, I, Pinho, D, Morato, M, Wilson, SP, Albino-Teixeira, A, Tavares, I Decrease in the Expression of N-Methyl-D-Aspartate Receptors in the Nucleus Tractus Solitarius Induces Antinociception and Increases Blood Pressure (2012) *JOURNAL OF NEUROSCIENCE RESEARCH* Volume: 90 Pages: 356-366

Marques-Lopes, J, Pinho, D, Albino-Teixeira, A, Tavares, I The hyperalgesic effects induced by the injection of angiotensin II into the caudal ventrolateral medulla are mediated by the pontine A(5) noradrenergic cell group (2010) *Brain Research* Vol 1325 Pages: 41-52

Marques-Lopes, J, Pinto, M, Pinho, D, Morato, M, Patinha, , Albino-Teixeira A, Tavares, I Microinjection of angiotensin II in the caudal ventrolateral medulla induces hyperalgesia (2009) *Neuroscience* Volume: 158 Pages: 1301-1310

Martins, I, Cabral, Pinto, A, Wilson, SP, Lima, D, Tavares, I Reversal of inflammatory pain by HSV-1-mediated overexpression of enkephalin in the caudal ventrolateral medulla

(2011) *European Journal of Pain* Volume: 15 Pages: 1008-1014

Martins, I, Costa-Araujo, S, Fadel, J, Wilson, SP, Lima, D, Tavares, I Reversal of neuropathic pain by HSV-1-mediated decrease of noradrenaline in a pain facilitatory area of the brain (2010) *PAIN* Volume: 151 Pages: 137-145

Martins, I, Pinto, , Wilson, SP, Lima, D, Tavares, I Dynamic of HSV-1 from a medullary pronociceptive centre: antinociception by overexpression of the preproenkephalin transgene (2008) *European Journal of Neuroscience* Volume: 28 Pages: 2075-2083

Martins, I; de Vries, MG; Teixeira-Pinto, A; Fadel, J; Wilson, S; Westerink, BHC; Tavares, I. Noradrenaline increases pain facilitation from the brain during inflammatory pain (2013) *Neuropharmacology* Volume: 71 Pages: 299-307

Martins-Oliveira M, Akerman S, Tavares I, Goadsby PJ. Neuropeptide Y inhibits the trigeminovascular pathway through NPY1 receptor: implications for migraine. *Pain* (2016), 157:1666-1673

Martins-Oliveira M, Akerman S, Holland FR, Hoffmann JR, Tavares I, Goadsby PJ. Neuroendocrine signaling modulates specific neural networks relevant to migraine, *Neurobiology of Disease*, May 2017, 101: 16-26.

Melnick IV, **Santos SFA & Safronov BV.** (2004) Mechanism of spike frequency adaptation in substantia gelatinosa neurons of rat. *J Physiol* 559: 383-395.

Melnick IV, **Santos SFA, Szokol K, Szucs P & Safronov BV.** (2004) Ionic basics of tonic firing in spinal substantia gelatinosa neurons of rat. *J Neurophysiol* 91: 646-655.

Melo A, Kokras N, Dalla C, Ferreira C, Ventura-Silva AP, Sousa N, et al. The positive effect on ketamine as a priming adjuvant in antidepressant treatment. *Translational psychiatry*. 2015;5:e573.

Melo A, Tavares I, Sousa N, Pego JM. Can the dopaminergic-related effects of general anesthetics be linked to mechanisms involved in drug abuse and addiction? *Acta anaesthesiologica Scandinavica*. 2015;59(7):822-9.

Mobina A, Cristiana Gaitero, Carlos Alexandre Ribeiro, Luís Miguel Santos, João Rodrigues Gomes, Sandra Marisa Oliveira, Pierre-Olivier Couraud, Babette Weksler, Ignacio Romero, Maria João Saraiva & Isabel Cardoso. Transthyretin participates in betaamyloid transport from the brain to the liver- involvement of the low density lipoprotein receptor-related protein 1? *Scientific Reports* | 6:20164 | DOI: 10.1038/srep20164.

Mobina A, Sara Costa Silva, Isabel Santana, Isabel Cardoso. Transthyretin stability is critical in assisting beta amyloid clearance - Relevance of Transthyretin stabilization in Alzheimer's Disease. *CNS Neuroscience & Therapeutics* (in press)

Monteiro C, Rebelo S, Galhardo V, Reguenga C, Lima D (2011) Postnatal expression of the homeobox gene *Prrxl1* (*Drg11*) is increased in inflammatory but not neuropathic pain. *European Journal of Pain* 15:477-481.

Monteiro C., Dourado M., Matos M., Duarte I., Lamas S., Galhardo V., Lima D. Critical Care and Survival of Fragile Animals: The case of *PRRXL1* Knockout Mice. *Applied Animal Behaviour Science* (2014) *APPLIED ANIMAL BEHAVIOUR SCIENCE* 158: 86-94

Monteiro C, Cardoso-Cruz H, Matos M, Dourado SM, Lima D, Galhardo V. Increased fronto-hippocampal connectivity in the *Prrxl1* knockout mouse model of congenital hypoalgesia. *Pain* 2016.

Monteiro CB, Costa MF, Reguenga C, Lima D, Castro DS, Monteiro FA. Paired related homeobox protein-like 1(*Prrxl1*) controls its own expression by a transcription autorepression mechanisms. *FEBS Letters* 588 (2014): 3475-3482

Monteiro CB, Midao L, Rebelo S, Reguenga C, Lima D, Monteiro FA. Zinc finger transcription factor *Cas21* expression is regulated by homeodomain transcription factor *Prrxl1* in embryonic spinal dorsal horn late born excitatory interneurons. *European Journal of Neuroscience* (2016): 1-11, doi:10.11/ejn.13214

Morgado C, Pereira-Terra P, Cruz CD, Tavares I. Minocycline completely reverses mechanical hyperalgesia in diabetic rats through microglia-induced changes in the expression of the potassium chloride co-transporter 2 (*KCC2*) at the spinal cord. *Diabetes Obes Metab.* 2011; 13:150-159.

Morgado C, Pereira-Terra P, Tavares I. Alpha-lipoic acid normalizes nociceptive neuronal activity at the spinal cord of diabetic rats. *Diabetes Obes Metab.* 2011, 13: 736-741.

Morgado C, Pinto-Ribeiro F, Tavares I. Diabetes affects the expression of GABA and potassium chloride cotransporter in the spinal cord: a study in streptozotocin diabetic rats. *Neurosci Lett.* 2008; 438:102-106.

Morgado C, Silva L, Pereira-Terra P, Tavares I. Changes in serotonergic and noradrenergic descending pain pathways during painful diabetic neuropathy: the preventive action of IGF1. *Neurobiol Dis.* 2011, 43: 275-284.

Morgado C, Tavares I. C-fos expression at the spinal dorsal horn of streptozotocin-induced diabetic rats. *Diabetes Metab Res Rev.* 2007; 23:644- 652.

Morgado C, Terra PP, Tavares I. Neuronal hyperactivity at the spinal cord and periaqueductal grey during painful diabetic neuropathy: effects of gabapentin. *Eur J Pain.* 2010; 14:693-699.

Nascimento D, Castro Lopes JM, Neto F. Satellite glial cells surrounding primary afferent neurons are activated and proliferate during monoarthritis in rats: is there a role for ATF3? *PLoS One* 2014, 23;9 (9): e108152

Nascimento D, Pozza D, Castro Lopes JM, Neto F. Neuronal injury marker ATF-3 is induced in primary afferent neurons of monoarthritic rats. *Neurosignals* 2011,19 (4):210-221

Nascimento, D.S, Potes, C.S., Soares, M.L., Ferreira, A.C., Malcangio, M, Castro-Lopes, J.M., Neto, F.L. "Drug-induced HSP90 inhibition alleviates pain in monoarthritic rats and alters the expression of new putative pain players at the DRG". *Molecular Neurobiology*, 2017 (in press)

Neto E, Cecília J. Alves, Daniela M. Sousa, Inês S. Alencastre, Ana H. Lourenço, Luís Leitão, Hyun R. Ryu, Noo L. Jeon, Rui Fernandes, Paulo Aguiar, Ramiro D. Almeida and Meriem Lamghari. Sensory neurons and osteoblasts: close partners in a microfluidic platform. *Integr. Biol.*, 2014, 6, 586

Neto E, Luís Leitão, Daniela M. Sousa,XCecília J. Alves,XIneês S. lencastre,XPaulo

Aguiar, and Meriem Lamghari. Compartmentalized Microfluidic Platforms: The Unrivaled Breakthrough of *In Vitro* Tools for Neurobiological Research. *The Journal of Neuroscience*, 2016 • 36(46):11573–11584

Neto, E, Alves, CJ, Leitao, L, et al. Axonal outgrowth, neuropeptides expression and receptors tyrosine kinase phosphorylation in 3D organotypic cultures of adult dorsal root ganglia, *PLoS ONE*, 2017.

Pereira S, Fontes F, Sonin T, Dias T, Fragoso M, Castro-Lopes JM, Lunet N. Neurological complications of breast cancer: A prospective cohort study. *The Breast* 24 (2015) 582-587

Pereira S, Fontes F, Sonin T, Dias T, Fragoso M, Castro-Lopes JM, Lunet N. Chemotherapy-induced peripheral neuropathy after neoadjuvant or adjuvant treatment of breast cancer: a prospective cohort study. *Support Care Cancer* DOI 10.1007/s00520-015-2935-y

Pereira S, Fontes F, Sonin T, Dias T, Fragoso M, Castro-Lopes JM, Lunet N. Neurological complications of breast cancer: study protocol of a prospective cohort study. *BMJ open*, 2014; 4(10):e006301

Pereira S, Fontes F, Sonin T, Dias T, Fragoso M, Castro-Lopes JM, Lunet N. Neuropathic pain after breast cancer treatment: characterization and risk factors (2017). *J Pain Symptom Manage*. 2017 Dec;54(6):877-888. doi: 10.1016/j.jpainsymman.2017.04.011.

Pinho R, Leonor C. Guedes, Lilach Soreq, Patricia P. Lobo, Tiago Mestre, Miguel Coelho, Mário M. Rosa, Nilza Gonçalves, Pauline Wales, Tiago Mendes, Gabriel Miltenberger-Miltényi, Fran Borovecki, Hermona Soreq, Joaquim J. Ferreira, Tiago F. Outeiro, Gene Expression Differences in Peripheral Blood of Parkinson's Disease Patients with Distinct Progression Profiles. *PLOS ONE* | DOI:10.1371/journal.pone.0157852 June 20, 2016:1-19

Paiva I, Pinho R, Pavlou MAS, Hennion M, Schuetz A-L, Rajput A, Szegő É, Kerimoglu C, Gerhardt E, Rego AC, Fischer A, Bonn S, Outeiro T. SODIUM BUTYRATE RESCUES DOPAMINERGIC CELLS FROM ALPHA-SYNUCLEIN-INDUCED TRANSCRIPTIONAL Deregulation and DNA Damage. *Human Molecular Genetics* (in press)

Pinto R, Frias B, Allen S, Dawbarn D, McMahon SB, Cruz F, et al. SEQUESTRATION OF BRAIN DERIVED NERVE FACTOR BY INTRAVENOUS DELIVERY OF TrkB-Ig(2) REDUCES BLADDER OVERACTIVITY AND NOXIOUS INPUT IN ANIMALS WITH CHRONIC CYSTITIS. *Neuroscience*. 2010;166(3):907-16.

Rebelo, S, Chen, ZF, Anderson, DJ, Lima, D Involvement of DRG11 in the development of the primary afferent nociceptive system (2006) *MOLECULAR AND CELLULAR NEUROSCIENCE* Volume: 33 Pages: 236-246

Rebelo, S, Lopes, C, Lima, D, Reguenga, C Expression of a Prrxl1 alternative splice variant during the development of the mouse nociceptive system (2009) *International Journal of Development Biology* Volume: 53 Issue: 7

Rebelo, S, Reguenga, C, Lopes, C, Lima, D Prrxl1 is Required for the Generation of a Subset of Nociceptive Glutamatergic Superficial Spinal Dorsal Horn Neurons (2010) *Development Dynamics* Volume: 239 Pages: 1684-1694

Rebelo, S, Reguenga, C, Osorio, L, Pereira, C, Lopes, C, Lima, D DRG11

immunohistochemical expression during embryonic development in the mouse (2007) *Development Dynamics* Volume: 236 Pages: 2653-2660

Rebouças ECC, Leal S, Sá SI. Regulation of NPY and α -MSH expression by estradiol in arcuate nucleus of Wistar female rats: a stereological study. *Neurological Research* 2016, 41-48

Rebouças ECC, Leal S, Silva SM, Sá SI. Changes in the female arcuate nucleus morphology and neurochemistry after chronic ethanol consumption and long-term withdrawal. *J. Chemical Neuroanatomy*, 77:30-40, 2016

Regadas I, Matos MR, Monteiro FA, Gómez-Skarmeta JL, Lima D, Bessa J, Casares F, Reguenga C. (2013). Several Cis-regulatory Elements Control mRNA Stability, Translation Efficiency, and Expression Pattern of Prrxl1 (Paired Related Homeobox Protein-like 1). *J Biol Chem*. Dec 20;288(51):36285-301. PMID: 24214975.

Regadas I, Soares-Dos-Reis R, Falcão M, Matos MR, Monteiro FA, Lima D, Reguenga C. (2014). Dual role of Tlx3 as modulator of Prrxl1 transcription and phosphorylation. *Biochim Biophys Acta – Gene Regulatory Mechanisms*. 1839(11):1121-1131. PubMed PMID: 25138281. (IF=6.332)

Sa, S, Madeira, MD Neuronal organelles and nuclear pores of hypothalamic ventromedial neurons are sexually dimorphic and change during the estrus cycle in the rat (2005) *NEUROSCIENCE* Volume: 133 Pages: 919-924

Sa, SI, Lukoyanova, E, Madeira, MD. Neuroscience effects of estrogens and progesterone on the synaptic organization of the hypothalamic ventromedial nucleus (2009) Volume: 162 Pages: 307-316.

Sa, SI, Madeira, MD Estrogen modulates the sexually dimorphic synaptic connectivity of the ventromedial nucleus (2005) *Journal of Comparative Neurology* Volume: 484 Pages: 68-79

Sa, SI, Pereira, PA, Paula-Barbosa, MM, Madeira, MD Role of neural afferents as mediators of estrogen effects on the hypothalamic ventromedial nucleus (2010) *Brain Research* Volume: 1366 Pages: 60-70

Santos SFA, Luz LL, Szucs P, Lima D, Derkach VA & Safronov BV. (2009) Transmission efficacy and plasticity in glutamatergic synapses formed by excitatory interneurons of the substantia gelatinosa in the rat spinal cord. *PlosOne* 4: e8047, p1-18.

Santos SFA, Melnick IV & Safronov BV. (2004) Selective postsynaptic inhibition of tonic-firing neurons in substantia gelatinosa by μ -opioid agonist. *Anesthesiology* 101: 1177-1183.

Santos SFA, Rebelo S, Safronov BV & Derkach VA. (2007) Excitatory interneurons dominate sensory processing in the spinal substantia gelatinosa of rat. *J Physiol* 581: 241- 254.

Seixas D, Foley P, Palace J, Lima D, Ramos I, Tracey I. Pain in multiple sclerosis: a systematic review of neuroimaging studies. *NeuroImage: Clinical* (in press)

Seixas D, Sá MJ, Galhardo V, Guimarães J, Lima D. Pain in Portuguese patients with multiple sclerosis. *Front Neurol* 2011; 2: 20;

Seixas, D, Lima, D Accuracy, reliability, validity and limitations of functional and structural magnetic resonance imaging data (2011) *CORTEX* Volume: 47; Pages: 1266-1269.

Seixas D, Jacqueline, Tracey I. Chronic pain disrupts the reward circuitry in Multiple sclerosis. *Eur. J. Neurosc.* 44: 192-198.

Silva, A, Cardoso-Cruz, H, Silva, F, Galhardo, V, Antunes, L Comparison of Anesthetic Depth Indexes Based on Thalamocortical Local Field Potentials in Rats (2010) *ANESTHESIOLOGY* Volume: 112 Pages: 355-363

Silva M, Amorim D, Almeida A, Tavares I, Pinto-Ribeiro F, Morgado C. Pronociceptive changes in the activity of rostroventromedial medulla (RVM) pain modulatory cells in the streptozotocin-diabetic rat. *Brain Research Bulletin.* 2013;96:39-44.

Silva M, Martins D, Charrua A, Piscitelli F, Tavares I, Morgado C, et al. Endovanilloid control of pain modulation by the rostroventromedial medulla in an animal model of diabetic neuropathy. *Neuropharmacology.* 2016;107:49-57

Silva, SM, Madeira, MD Effects of chronic alcohol consumption and withdrawal on the response of the male and female hypothalamic-pituitary-adrenal axis to acute immune stress (2012) *BRAIN RESEARCH* Volume: 1444 Pages: 27-37

Silva, SM, Madeira, MD, Ruela, C, Paula-Barbosa, MM Prolonged alcohol intake leads to irreversible loss of vasopressin and oxytocin neurons in the paraventricular nucleus of the hypothalamus (2002) *BRAIN RESEARCH* Volume: 925 Pages: 76-88

Silva, SM, Paula-Barbosa, MM, Madeira, MD, Prolonged alcohol intake leads to reversible depression of corticotropin-releasing hormone and vasopressin immunoreactivity and mRNA levels in the parvocellular neurons of the paraventricular nucleus (2002) *BRAIN RESEARCH*; Volume: 954; Pages: 82-93

Silva, SM, Santos-Marques, MJ, Madeira, D Sexually dimorphic response of the hypothalamo-pituitary-adrenal axis to chronic alcohol consumption and withdrawal (2009) *BRAIN RESEARCH* Volume: 1303 Pages: 61-73

Sousa M, Aguiar P. Building, simulating and visualizing large spiking neural networks with neuralSyms. *Neurocomputing* 123: 372-380 (2014)

Sousa M, Szucs P, Lima D, Aguiar P. The pronociceptive dorsal reticular nucleus contains mostly tonic neurons and shows a high prevalence of spontaneous activity in block preparation. *J. Neurophysiol* 11:1507-1518 (2014)

Szucs P, Luz LL, Lima D & Safronov BV. Local axon collaterals of lamina I projection neurons in the spinal cord of young rats. *J Comp Neurol* 518, 2645-2665, 2010.

Szucs P, Luz LL, Pinho R, Aguiar P, Antal Zs, Tiong SY, Todd A, Safronov BV. Axon Diversity of Lamina I Local-Circuit Neurons in the Lumbar Spinal Cord *J Comp Neurol* 521: 2719-2741 (2013) DOI: 10.1002/cne.23311

Vaz RP, Pedro A. Pereira and M. Dulce Madeira. Age Effects on the Nucleus of the Lateral Olfactory Tract of the Rat. *The Journal of Comparative Neurology* 524:759-771

(2016)

Vaz RP, Armando Cardoso, Susana I. Sa´, Pedro A. Pereira¹, M. Dulce Madeira. The integrity of the nucleus of the lateral olfactory tract is essential for the normal functioning of the olfactory system. *Brain Struct Funct* DOI 10.1007/s00429-017-1422-2 (2017)

Vieira JB, Almeida PR, Ferreira-Santos F, Barbosa F, Marques-Teixeira J, Marsh AA. Distinct neural activation patterns underlie economic decisions in high and low psychopathy scorers. *Social cognitive and affective neuroscience*. 2013 Jul 16.

Vieira, JB, Marsh, AA Don't stand so close to me: psychopathy and the regulation of interpersonal distance. *FRONTIERS IN HUMAN NEUROSCIENCE* 7: Art.no. 907 (2014) DOI: 10.3389/fnhum.2013.00907.

Vieira JB, Ferreira-Santos F, Almeida PR, Barbosa F, Marques-Teixeira, and Marsh AA. Psychopathic traits are associated with cortical and subcortical volume alterations in healthy individuals (2015), *Social Cognitive and Affective Neuroscience*, 10(12):1693-704.

Xu Y, **Lopes C**, Wende H, Guo Z, Cheng L, Birchmeier C, Ma Q. (2013) Ontogeny of excitatory spinal neurons processing distinct somatic sensory modalities. *J Neurosci*. 2013 Sep 11;33(37):14738-48.